


PRODUCT INFORMATION  
**IL-34 (Interleukin-34), Human**

<b>Catalog number</b>	C01040-5UG / C01040-20UG / C01040-100UG
<b>Package</b>	5 µg / 20 µg / 100 µg
<b>Description</b>	Interleukin-34 (IL-34) is a protein that promotes the proliferation, survival and differentiation of monocytes and macrophages. It also promotes the release of proinflammatory chemokines, and thereby plays an important role in innate immunity and in inflammatory processes. IL-34 plays an important role in the regulation of osteoclast proliferation and differentiation, and in the regulation of bone resorption. Signaling via CSF1R and its downstream effectors stimulates phosphorylation of MAPK1/ERK2 AND MAPK3/ERK1.
<b>Source</b>	<i>Escherichia coli</i>
<b>Sequence</b>	MNEPLEMWPLTQNEECTVTGFLRDKLQYRSRLQYMKHYFPINYKISVPYEGVF RIANVTRLQRAQVSERELRYLWVLVLSLATESVQDVLLEGHPSWKYLQEVETLL LNVQQGLTDVEVSPKVESVLSLLNAPGPNLKLVRPKALLDNCFRVMELLYCSCC KQSSVLNWQDCEVSPQSCSPEPSLQYAATQLYPPPPWSPSSPHSTGSRP VRAQGEGLLP with polyhistidine tag at the C-terminus
<b>Endotoxin level</b>	<0.1 EU per 1 µg of the protein by the LAL method.
<b>Activity</b>	Measure by its ability to induce human peripheral blood monocytes (PBMCs) proliferation. The ED <sub>50</sub> for this effect is <30 pg/mL.
<b>Purity</b>	>98% as determined by SDS-PAGE. Ni-NTA chromatography
<b>Formulation</b>	The protein was lyophilized from a solution containing 1X PBS, pH 8.0.
<b>Reconstitution</b>	It is recommended to reconstitute the lyophilized protein in sterile H <sub>2</sub> O to a concentration not less than 100 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.
<b>Storage</b>	Lyophilized protein should be stored at -20°C. Upon reconstitution, protein aliquots should be stored at -20°C or -80°C.
<b>Note</b>	Please use within one month after protein reconstitution.



SDS-PAGE analysis of recombinant human IL-34

For Research Use Only.