


## PRODUCT INFORMATION

**FGF-5 (Fibroblast growth factor-5), Human**

<b>Catalog number</b>	C01095-5UG / C01095-20UG / C01095-100UG
<b>Package</b>	5 µg / 20 µg / 100 µg
<b>Description</b>	FGF-5 is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth, and invasion. In the nervous system, FGF-5 has been most often identified in neurons associated with the limbic system, notably in neurons of the olfactory bulb and pyramidal cells of the hippocampus. Hippocampal FGF-5 is suggested to serve as a neurotrophic and differentiative factor for cholinergic and serotonergic neurons projecting to this region.
<b>Source</b>	<i>Escherichia coli</i>
<b>Sequence</b>	MAWAHGEKRLAPKGQPGPAATDRNPIGSSSRQSSSSAMSSSSASSSPAASLG SQGSGLEQSSFQWSPSGRRTGSLYCRVGIGFHLQIYDPDGKVNGSHEANMLSV LEIFAVSQGIVGIRGVFSNKFLAMSKKGLHASAKFTDDCKFRERFQENSYNTY ASAIHRTEKTGREWYVALNKRKAKRGCSRPVKPQHISTHFLPRFKQSEQPEL SFTVTVPEKKNPPSPIKSK IPLSAPRKNT NSVKYRLKFR FG with polyhistidine tag at the C-terminus
<b>Endotoxin level</b>	<0.1 EU per 1 µg of the protein by the LAL method.
<b>Activity</b>	Measure by its ability to induce 3T3 cells proliferation. The ED <sub>50</sub> for this effect is <0.7 ng/mL. The specific activity of recombinant human FGF-5 is >1.4 x 10 <sup>6</sup> IU/mg.
<b>Purity</b>	>95% as determined by SDS-PAGE. Ni-NTA chromatography
<b>Formulation</b>	The protein was lyophilized from a solution containing 1X PBS, pH 8.0.
<b>Reconstitution</b>	It is recommended to reconstitute the lyophilized protein in sterile H <sub>2</sub> O to a concentration not less than 100 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.
<b>Storage</b>	Lyophilized protein should be stored at -20°C. Upon reconstitution, protein aliquots should be stored at -20°C or -80°C.
<b>Note</b>	Please use within one month after protein reconstitution.



SDS-PAGE analysis of recombinant human FGF-5

For Research Use Only.