
## **PRODUCT INFORMATION**

## FGF-11 isoform 1 (Fibroblast growth factor-11 isoform 1), Human

Package 5 μg / 20 μg / 100 μg   FGF-11 is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. The expression pattern of the mouse homolog implies a role in nervous system development.   Source Escherichia coli   MAALASSLIRQKREVREPGGSRPVSAQRRVCPRGTKSLCQKQLLILLSKVRLCG GRPARPDRGPEPQLKGIVTKLFCRQGFYLQANPDGSIQGTPEDTSSFTHFNLIP   VGLRVVTIQSAKLGHYMAMNAEGLLYSSPHFTAECRFKECVFENYYVLYASALY RQRRSGRAWYLGLDKEGQVMKGNRVKKTKAAAHFLPKLLEVAMYQEPSLHSV PEASPSSPPAP with polyhistidine tag at the C-terminus   Endotoxin level <0.01 EU per 1 μg of the protein by the LAL method.   Activity Measure by its ability to induce 3T3 cells proliferation. The ED <sub>50</sub> for this effect is <0.2 ng/mL.		
FGF-11 is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. The expression pattern of the mouse homolog implies a role in nervous system development.   Source Escherichia coli   MAALASSLIRQKREVREPGGSRPVSAQRRVCPRGTKSLCQKQLLILLSKVRLCG GRPARPDRGPEPQLKGIVTKLFCRQGFYLQANPDGSIQGTPEDTSSFTHFNLIP   VGLRVVTIQSAKLGHYMAMNAEGLLYSSPHFTAECRFKECVFENYYVLYASALY RQRRSGRAWYLGLDKEGQVMKGNRVKKTKAAAHFLPKLLEVAMYQEPSLHSV PEASPSSPPAP with polyhistidine tag at the C-terminus   Endotoxin level <0.01 EU per 1 µg of the protein by the LAL method.   Activity Measure by its ability to induce 3T3 cells proliferation. The ED <sub>50</sub> for this effect is <0.2 ng/mL.	Catalog number	C01101-5UG / C01101-20UG / C01101-100UG
Descriptionmembers possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. The expression pattern of the mouse homolog implies a role in nervous system development.SourceEscherichia coliSequenceMAALASSLIRQKREVREPGGSRPVSAQRRVCPRGTKSLCQKQLLILLSKVRLCG GRPARPDRGPEPQLKGIVTKLFCRQGFYLQANPDGSIQGTPEDTSSFTHFNLIP VGLRVVTIQSAKLGHYMAMNAEGLLYSSPHFTAECRFKECVFENYYVLYASALY RQRRSGRAWYLGLDKEGQVMKGNRVKKTKAAAHFLPKLLEVAMYQEPSLHSV PEASPSSPPAP with polyhistidine tag at the C-terminusEndotoxin level<0.01 EU per 1 µg of the protein by the LAL method.	Package	5 µg / 20 µg / 100 µg
MAALASSLIRQKREVREPGGSRPVSAQRRVCPRGTKSLCQKQLLILLSKVRLCG GRPARPDRGPEPQLKGIVTKLFCRQGFYLQANPDGSIQGTPEDTSSFTHFNLIP VGLRVVTIQSAKLGHYMAMNAEGLLYSSPHFTAECRFKECVFENYYVLYASALY RQRRSGRAWYLGLDKEGQVMKGNRVKKTKAAAHFLPKLLEVAMYQEPSLHSV PEASPSSPPAP with polyhistidine tag at the C-terminusEndotoxin level<0.01 EU per 1 µg of the protein by the LAL method.ActivityMeasure by its ability to induce 3T3 cells proliferation. The ED <sub>50</sub> for this effect is <0.2 ng/mL.	Description	members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. The expression pattern
SequenceGRPARPDRGPEPQLKGIVTKLFCRQGFYLQANPDGSIQGTPEDTSSFTHFNLIP VGLRVVTIQSAKLGHYMAMNAEGLLYSSPHFTAECRFKECVFENYYVLYASALY RQRRSGRAWYLGLDKEGQVMKGNRVKKTKAAAHFLPKLLEVAMYQEPSLHSV PEASPSSPPAP with polyhistidine tag at the C-terminusEndotoxin level<0.01 EU per 1 µg of the protein by the LAL method.	Source	Escherichia coli
Activity Measure by its ability to induce 3T3 cells proliferation. The ED <sub>50</sub> for this effect is <0.2 ng/mL.	Sequence	GRPARPDRGPEPQLKGIVTKLFCRQGFYLQANPDGSIQGTPEDTSSFTHFNLIP VGLRVVTIQSAKLGHYMAMNAEGLLYSSPHFTAECRFKECVFENYYVLYASALY RQRRSGRAWYLGLDKEGQVMKGNRVKKTKAAAHFLPKLLEVAMYQEPSLHSV
Activity <0.2 ng/mL.	Endotoxin level	<0.01 EU per 1 $\mu$ g of the protein by the LAL method.
Purity >98% as determined by SDS-PAGE. Ni-NTA chromatography	Activity	
	Purity	>98% as determined by SDS-PAGE. Ni-NTA chromatography
<b>Formulation</b> The protein was lyophilized from a solution containing 1X PBS, pH 7.4.	Formulation	The protein was lyophilized from a solution containing 1X PBS, pH 7.4.
ReconstitutionIt is recommended to reconstitute the lyophilized protein in sterile H₂O to a concentration not less than 100 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.	Reconstitution	concentration not less than 100 $\mu$ g/mL and incubate the stock solution for at least
Storage Lyophilized protein should be stored at -20°C. Upon reconstitution, protein aliquots should be stored at -20°C or -80°C.	Storage	
<b>Note</b> Please use within one month after protein reconstitution.	Note	Please use within one month after protein reconstitution.





For Research Use Only.

