


PRODUCT INFORMATION

Annexin V, Human

Catalog number	C01168-5UG / C01168-20UG / C01168-100UG
Package	5 µg / 20 µg / 100 µg
Description	Annexin V is a cellular protein in the annexin group. In flow experiment, annexin V is commonly used to detect apoptotic cells by its ability to bind to phosphatidylserine, a marker of apoptosis when it is on the outer leaflet of the plasma membrane. Annexin V has been proposed to play a role in the inhibition of blood coagulation by competing for phosphatidylserine binding sites with prothrombin and also to inhibit the activity of phospholipase A1.
Source	<i>Escherichia coli</i>
Sequence	MAQVLRGTVTDFPGFDERADAETLRKAMKGLGTDEESILTLTSSRSNAQRQEIS AAFKTLFGRDLLDDLKSELTGKFEKLIVALMKPSRLYDAYELKHALKGAGTNEKV LTEIIASRTPEELRAIKQVYEEYEGSSLEDDVVGDTSGYYQRMLVLLQANRDPD AGIDEAQVEQDAQALFQAGELKWGTDEEKFITIFGTRSVSHLRKVFVKYMTISGF QIETIDRETSGNLEQLLLAVVKSIRSIPAYLAETLYYAMKGAGTDDHTLIRVMVS RSEIDLFNIRKEFRKNFATSLYSMIKGDTSGDYKKALLLLCGEDD with polyhistidine tag at the C-terminus
Endotoxin level	<0.1 EU per 1 µg of the protein by the LAL method.
Purity	>98% as determined by SDS-PAGE. Ni-NTA chromatography
Formulation	The protein was lyophilized from a solution containing 1X PBS, pH 7.4.
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile H ₂ O to a concentration not less than 100 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.
Storage	Lyophilized protein should be stored at -20°C. Upon reconstitution, protein aliquots should be stored at -20°C or -80°C.
Note	Please use within one month after protein reconstitution.



SDS-PAGE analysis of recombinant human Annexin V

For Research Use Only.