


PRODUCT INFORMATION  
**IL-5 (Interleukin-5), Mouse**

<b>Catalog number</b>	C02007-5UG / C02007-20UG / C02007-100UG
<b>Package</b>	5 µg / 20 µg / 100 µg
<b>Description</b>	Interleukin 5 (IL-5) is an interleukin produced by type-2 T helper cells and mast cells. Through binding to the interleukin-5 receptor, interleukin 5 stimulates B cell growth and increases immunoglobulin secretion - primarily IgA. It is also a key mediator in eosinophil activation.
<b>Source</b>	<i>Escherichia coli</i>
<b>Sequence</b>	MEIPMSTVVKETLTQLSAHRALLTSNETMRLPVPTHKNHQLCIGEIFQGLDILKN QTVRGGTVEMLFQNLSLIKKYIDRQKEKCGEERRRTRQFLDYLQEFLGVMSTE WAMEG with polyhistidine tag at the C-terminus
<b>Endotoxin level</b>	<0.1 EU per 1 µg of the protein by the LAL method.
<b>Activity</b>	Measure by its ability to induce TF-1 cells proliferation. The ED <sub>50</sub> for this effect is <0.2 ng/mL. The specific activity of recombinant mouse IL-5 is > 5 x 10 <sup>6</sup> IU/mg.
<b>Purity</b>	>98% as determined by SDS-PAGE. Ni-NTA chromatography
<b>Formulation</b>	The protein was lyophilized from a solution containing 1X PBS, pH 7.4.
<b>Reconstitution</b>	It is recommended to reconstitute the lyophilized protein in sterile H <sub>2</sub> O to a concentration not less than 100 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.
<b>Storage</b>	Lyophilized protein should be stored at -20°C. Upon reconstitution, protein aliquots should be stored at -20°C or -80°C.
<b>Note</b>	Please use within one month after protein reconstitution.



SDS-PAGE analysis of recombinant mouse IL-5

For Research Use Only.