


PRODUCT INFORMATION
IL-17F (Interleukin-17F), Mouse

Catalog number	C02021-5UG / C02021-20UG / C02021-100UG
Package	5 µg / 20 µg / 100 µg
Description	The protein encoded by this gene is a cytokine that shares sequence similarity with IL17. This cytokine is expressed by activated T cells, and has been shown to stimulate the production of several other cytokines, including IL6, IL8, and CSF2/GM-CSF. This cytokine is also found to inhibit the angiogenesis of endothelial cells and induce endothelial cells to produce IL2, TGFB1/TGFB, and monocyte chemoattractant protein-1.
Source	<i>Escherichia coli</i>
Sequence	MRKNPKAGVPALQKAGNCPPLEDNTVRVDIRIFNQNQGISVPREFQNRSSSPW DYNITRDPHRFPSEIAEAQCRHSGCINAQQQEDSTMNSVAIQQEILVLRREPQG CSNSFRLEKMLLKVGCTCVKPIVHQAA with polyhistidine tag at the C-terminus
Endotoxin level	<0.1 EU per 1 µg of the protein by the LAL method.
Activity	Measure by its ability to induce IL-6 secretion in 3T3 cells. The ED ₅₀ for this effect is <100 ng/mL.
Purity	>98% as determined by SDS-PAGE. Ni-NTA chromatography
Formulation	The protein was lyophilized from a solution containing 20 mM sodium citrate, pH 4.5.
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile H ₂ O to a concentration not less than 100 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.
Storage	Lyophilized protein should be stored at -20°C. Upon reconstitution, protein aliquots should be stored at -20°C or -80°C.
Note	Please use within one month after protein reconstitution.



SDS-PAGE analysis of recombinant mouse IL-17F

For Research Use Only.