
PRODUCT INFORMATION

IGF-II (Insulin-like growth factor-II), Mouse

-	
Catalog number	C02063-5UG / C02063-20UG / C02063-100UG
Package	5 μg / 20 μg / 100 μg
Description	Insulin-like growth factor 2 (IGF-2) is one of three protein hormones that share structural similarity to insulin. The major role of IGF-2 is as a growth promoting hormone during gestation. It is believed to be a major fetal growth factor in contrast to Insulin-like growth factor 1, which is a major growth factor in adults.
Source	Escherichia coli
Sequence	YGPGETLCGGELVDTLQFVCSDRGFYFSRPSSRANRRSRGIVEECCFRSCDLA LLETYCATPAKSE with polyhistidine tag at the N-terminus
Endotoxin level	<0.1 EU per 1 µg of the protein by the LAL method.
Activity	Measure by its ability to induce MCF-7 cells proliferation. The ED $_{50}$ for this effect is <6 ng/mL. The specific activity of recombinant mouse IGF-II is > 1.5 x 10^5 IU/mg.
Purity	>98% as determined by SDS-PAGE. Ni-NTA chromatography
Formulation	The protein was lyophilized from a solution containing 1X PBS, pH 8.0.
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile H_2O to a concentration not less than 100 μ g/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.
Storage	Lyophilized protein should be stored at -20°C. Upon reconstitution, protein aliquots should be stored at -20°C or -80°C.
Note	Please use within two weeks after protein reconstitution.



For Research Use Only.