


PRODUCT INFORMATION

Flt-3 Ligand (Fms-related tyrosine kinase-3 ligand), Swine

Catalog number	C03014-5UG / C03014-20UG / C03014-100UG
Package	5 µg / 20 µg / 100 µg
Description	Fms-related tyrosine kinase 3 ligand (FLT3LG) is a protein which in humans is encoded by the FLT3LG gene. FLT3 ligand is a receptor for the fl cytokine has a tyrosine-protein kinase activity & a growth factor that regulates proliferation of early hematopoietic cells. Flt3-Ligand synergizes with other CSFs and interleukins to induce growth and differentiation.
Source	<i>Escherichia coli</i>
Sequence	MSPDCSFPHSPISSTFANTIRQLSDYLLQDYPVTVASNLQDDELTCGAFWRLVLA QRWVGQLKTVAGSQMQKLLLEAVNTEIVFVTSCALQLPLPSCRFVQANISHLLQ DTSQQLVALKPWITRRNFSRCLELQCQPDPSTLLPPRSPGALEATSLPAPQASL LLLLLLLLLPAALLLL with polyhistidine tag at the C-terminus
Endotoxin level	<0.1 EU per 1 µg of the protein by the LAL method.
Activity	Measure by its ability to induce proliferation in OCI-AML5 cells. The ED ₅₀ for this effect is <5 ng/mL.
Purity	>95% as determined by SDS-PAGE. Ni-NTA chromatography
Formulation	The protein was lyophilized from a solution containing 1X PBS, pH 7.4.
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile H ₂ O to a concentration not less than 100 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.
Storage	Lyophilized protein should be stored at -20°C. Upon reconstitution, protein aliquots should be stored at -20°C or -80°C.
Note	Please use within one month after protein reconstitution.



SDS-PAGE analysis of recombinant swine Flt-3 Ligand

For Research Use Only.