


PRODUCT INFORMATION
CD27L (CD27 ligand), Mouse

Catalog number	C02045-5UG / C02045-20UG / C02045-100UG
Package	5 µg / 20 µg / 100 µg
Description	CD27 Ligand, which belongs to the TNF superfamily, is a type II transmembrane protein. Like NK cells, CD27L is also located in activated T and B lymphocytes. Along with NK enhancement, CD27L and its receptor regulate the immune response by raising expansion and differentiation of T cell. CD27 signaling can act like a co-stimulatory effector to sustain survival of the CD8+ T cells, mainly by increasing expression of the IL-2 gene.
Source	<i>Escherichia coli</i>
Sequence	QQQRLLLEHPEPHTAELQLNLTVPRKDPTLRWGAGPALGRSFTTHGPELEEGHLR IHQDGLYRLHIQVTLANCSSPGSTLQHRATLAVGICSPA AHGISLLRGRFGQDCT VALQRLTYLVHGDVLTNLTLPLPSRNADETFFGVQWICP with polyhistidine tag at the N-terminus
Endotoxin level	<0.1 EU per 1 µg of the protein by the LAL method.
Activity	Measure by its ability to induce proliferation in mouse T cells in the presence of the anti-CD3 antibody. The ED ₅₀ for this effect is <4.5 µg/mL.
Purity	>98% as determined by SDS-PAGE. Ni-NTA chromatography
Formulation	The protein was lyophilized from a solution containing 1X PBS, pH7.4.
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile H ₂ O to a concentration not less than 100 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.
Storage	Lyophilized protein should be stored at -20°C. Upon reconstitution, protein aliquots should be stored at -20°C or -80°C.
Note	Please use within one month after protein reconstitution.



SDS-PAGE analysis of recombinant mouse CD27L

For Research Use Only.