


## PRODUCT INFORMATION

**TRAIL (TNF-related apoptosis-inducing ligand), Mouse**

<b>Catalog number</b>	C02049-5UG / C02049-20UG / C02049-100UG
<b>Package</b>	5 µg / 20 µg / 100 µg
<b>Description</b>	TRAIL acts as a cytotoxic protein, through activating rapid apoptosis in tumor cells (not in normal cells). TRAIL conducts apoptosis through binding to DR4 and DR5, which are two death-signaling receptors belonging to the TNFR superfamily of transmembrane proteins. These receptor contain a cytoplasmic "death domain", which can initiate the cell's apoptotic process.
<b>Source</b>	<i>Escherichia coli</i>
<b>Sequence</b>	MPRGGRPQKVAAHITGITRRSNSALIPISKDGKTLGQKIESWESSRKGHSFLNH VLFRRNGELVIEQEGLYIYSQTYFRFQEAEDASKMVSKDKVRTKQLVQYIYKYTS YPDPIVLMKSARNSCWSRDAEYGLYSIQGGLFELKKNDRIFVSVTNEHLMDLD QEASFFGAFLIN with polyhistidine tag at the C-terminus
<b>Endotoxin level</b>	<0.1 EU per 1 µg of the protein by the LAL method.
<b>Activity</b>	Measure by its ability to induce cytotoxicity in L929 cells in the presence of actinomycin D. The ED <sub>50</sub> for this effect is <1 ng/mL.
<b>Purity</b>	>98% as determined by SDS-PAGE. Ni-NTA chromatography
<b>Formulation</b>	The protein was lyophilized from a solution containing 1X PBS, pH 7.4.
<b>Reconstitution</b>	It is recommended to reconstitute the lyophilized protein in sterile H <sub>2</sub> O to a concentration not less than 100 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.
<b>Storage</b>	Lyophilized protein should be stored at -20°C. Upon reconstitution, protein aliquots should be stored at -20°C or -80°C.
<b>Note</b>	Please use within one month after protein reconstitution.



SDS-PAGE analysis of recombinant mouse TRAIL

For Research Use Only.