

PRODUCT INFORMATION  
**CXCL3 (C-X-C motif chemokine 3), Mouse**

<b>Catalog number</b>	C02075-5UG / C02075-20UG / C02075-100UG
<b>Package</b>	5 µg / 20 µg / 100 µg
<b>Description</b>	CXCL3 is an ELR CXC chemokine. Its structural and functional characteristics are similar to GRO1 (CXCL1), GRO2 (CXCL2), and interleukin-8 (CXCL8). CXC chemokines is critical in the phase I inflammation, in which the PMN cells are rapidly chemoattracted. In the next phase of inflammation, the CC chemokines (MCPs) attract different cell subpopulations such as T cells, monocytes, basophils, and eosinophils. MMP12, primarily released by macrophages, can modulate the activity of ELR-CXC chemokines and cleavage human CXCL1, CXCL2 and CXCL3 within the ELR sequence at Glu6-Leu7.
<b>Source</b>	<i>Escherichia coli</i>
<b>Sequence</b>	AVVASELRCQCLNTLPRVDFETIQSLTVTPPGPHCTQTEVIATLKDGQEVCLNP QGPRLQIIKKILKSGKSS with polyhistidine tag at the N-terminus
<b>Endotoxin level</b>	<0.1 EU per 1 µg of the protein by the LAL method.
<b>Activity</b>	Measure by its ability to chemoattract BaF3 cells transfected with human CXCR2. The ED <sub>50</sub> for this effect is <80 ng/mL.
<b>Purity</b>	>98% as determined by SDS-PAGE. Ni-NTA chromatography
<b>Formulation</b>	The protein was lyophilized from a solution containing 1X PBS, pH 7.4.
<b>Reconstitution</b>	It is recommended to reconstitute the lyophilized protein in sterile H <sub>2</sub> O to a concentration not less than 100 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.
<b>Storage</b>	Lyophilized protein should be stored at -20°C. Upon reconstitution, protein aliquots should be stored at -20°C or -80°C.
<b>Note</b>	Please use within one month after protein reconstitution.



SDS-PAGE analysis of recombinant mouse CXCL3

For Research Use Only.