


PRODUCT INFORMATION

IFN gamma (Interferon gamma), Swine

Catalog number	C03012-5UG / C03012-20UG / C03012-100UG
Package	5 µg / 20 µg / 100 µg
Description	Interferon-γ is a effective multifunctional cytokine which is released mainly by activated NK cells and T cells. IFN-γ is primarily characterized based on its anti-viral activities, and then been proved to have several functions such as anti-proliferative, immune-regulatory, and pro-inflammatory activities. IFN-γ can upregulate expression of MHC class I and II antigen by antigen-presenting cells.
Source	<i>Escherichia coli</i>
Sequence	MQAPFFKEITILKDYFNASTSDVPNGGPLFLEILKNWKEESDKKIIQSQIVSFYFK FFEIFKDNQAIQRSMDVIKQDMFQRFLNGSSGKLNDFEKLKIPVDNLQIQRKAIS ELIKVMNDLSPRSNLRKRKRSQTMFQGQRASK with polyhistidine tag at the C-terminus
Endotoxin level	<0.01 EU per 1 µg of the protein by the LAL method.
Activity	Measure by its ability to protect PK15 cells infected with encephalomyocarditis (EMC) virus. The ED ₅₀ for this effect is <40 pg/mL.
Purity	>95% as determined by SDS-PAGE. Ni-NTA chromatography
Formulation	The protein was lyophilized from a solution containing 1X PBS, pH 7.4.
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile H ₂ O to a concentration not less than 100 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.
Storage	Lyophilized protein should be stored at -20°C. Upon reconstitution, protein aliquots should be stored at -20°C or -80°C.
Note	Please use within one month after protein reconstitution.



SDS-PAGE analysis of recombinant swine IFN gamma

For Research Use Only.