

PRODUCT INFORMATION

CCL4 (C-C motif chemokine ligand 4), Human

Catalog number	C01141-5UG / C01141-20UG / C01141-100UG
Package	5 µg / 20 µg / 100 µg
Description	CCL4 act as an important pro-inflammatory cytokine during acute and chronic inflammatory responses. CCL4 also play a role as an attractant NK cells, dendritic cells, monocytes, and lymphocytes to sites of injury and inflammation. As CCL4 conducts signaling through G-protein-coupled receptor CCR5, which is a major co-receptor for M-tropic HIV strains. Thus, binding of CCL4 to CCR5 inhibits HIV entry and reduces the cell surface expression of CCR5. CCL4 has been identified as one of the major anti-HIV factors produced by CD8+ T cells.
Source	<i>Escherichia coli</i>
Sequence	APMGSDPPTACCFSTARKLPHNFVVDYYETSSLCSQPAVVFQTKRGKQVCA DPSESWVQEYVYDLELN
Endotoxin level	<0.1 EU per 1 µg of the protein by the LAL method.
Activity	Measure by its ability to chemoattract BaF3 cells transfected with human CCR5. The ED ₅₀ for this effect is <10 ng/mL.
Purity	>98% as determined by SDS-PAGE. Ni-NTA chromatography
Formulation	The protein was lyophilized from a solution containing 50 mM Tris and 150 mM NaCl, pH 8.5.
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile H ₂ O to a concentration not less than 100 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.
Storage	Lyophilized protein should be stored at -20°C. Upon reconstitution, protein aliquots should be stored at -20°C or -80°C.
Note	Please use within one month after protein reconstitution.



SDS-PAGE analysis of recombinant human CCL4

For Research Use Only.