


PRODUCT INFORMATION
IL-3 (Interleukin-3), Mouse

Catalog number	C02005-5UG / C02005-20UG / C02005-100UG
Package	5 µg / 20 µg / 100 µg
Description	Interleukin 3 is an interleukin, a type of biological signal (cytokine) that can improve the body's natural response to disease as part of the immune system. It acts by binding to the interleukin-3 receptor. Interleukin 3 stimulates the differentiation of multipotent hematopoietic stem cells into myeloid progenitor cells or, with the addition of IL-7, into lymphoid progenitor cells. In addition, IL-3 stimulates proliferation of all cells in the myeloid lineage (granulocytes, monocytes, and dendritic cells), in conjunction with other cytokines, e.g., Erythropoietin (EPO), Granulocyte macrophage colony-stimulating factor (GM-CSF), and IL-6. It is secreted by basophils and activated T cells to support growth and differentiation of T cells from the bone marrow in an immune response.
Source	<i>Escherichia coli</i>
Sequence	MDTHRLTRLNCSSIVKEIIGKLPEPELKTDDDEGPSLRNKSFRRVNLKSFVESQG EVDPEDRYVIKSNLQKLNCLPTSANDSALPGVFIRDLDLDFRKKLRFYMVHLND LETVLTSRPPQPASGSVSPNRGTVEC with polyhistidine tag at the C-terminus
Endotoxin level	<0.1 EU per 1 µg of the protein by the LAL method.
Activity	Measure by its ability to induce NFS-60 cells proliferation. The ED ₅₀ for this effect is <85 pg/mL. The specific activity of recombinant mouse IL-3 is approximately >1x 10 ⁷ IU/mg.
Purity	≥95% as determined by SDS-PAGE and HPLC. Purified by Ni-NTA chromatography.
Formulation	The protein was lyophilized from a solution containing 1X PBS, pH 7.4.
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile H ₂ O to a concentration not less than 100 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.
Storage	Lyophilized protein should be stored at -20°C. Upon reconstitution, protein aliquots should be stored at -20°C or -80°C.
Note	Please use within one month after protein reconstitution.



SDS-PAGE analysis of recombinant mouse IL-3

For Research Use Only.