


PRODUCT INFORMATION

TGF beta 1 (Transforming growth factor beta 1), Swine

Catalog number	C03015-5UG / C03015-20UG / C03015-100UG
Package	5 µg / 20 µg / 100 µg
Description	Transforming growth factor beta 1 or TGF-β1 is a polypeptide member of the transforming growth factor beta superfamily of cytokines. It is a secreted protein that performs many cellular functions, including the control of cell growth, cell proliferation, cell differentiation, and apoptosis. In humans, TGF-β1 is encoded by the TGFB1 gene.
Source	<i>Escherichia coli</i>
Sequence	MALDTNYCFSSTEKNCCVRQLYIDFRKDLGWKWIHEPKGYHANFCLGPCPYIW SLDTQYSKVLALYNQHNP GASAAPCCVPQALEPLPIVYVGRKPKVEQLSNMIV RSCKCS with polyhistidine tag at the C-terminus
Endotoxin level	<0.1 EU per 1 µg of the protein by the LAL method.
Activity	Measure by its ability to inhibit IL-4-induce proliferation in HT-2 cells. The ED ₅₀ for this effect is <0.1 ng/mL.
Purity	>98% as determined by SDS-PAGE analysis. Purified by Ni-NTA chromatography
Formulation	The protein was lyophilized from a solution containing 20 mM sodium citrate, 0.2 M NaCl pH 4.5.
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile H ₂ O to a concentration not less than 100 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved. In some experiments, it recommends to add 10 mM HCl when reconstitute lyophilized protein.
Storage	Lyophilized protein should be stored at -20°C. Upon reconstitution, protein aliquots should be stored at -20°C or -80°C.
Note	Please use within one month after protein reconstitution.



SDS-PAGE analysis of recombinant swine TGF beta 1

For Research Use Only.