


PRODUCT INFORMATION

CD40L (CD40 ligand), Human

v. 231001

Catalog number	C01054-5UG / C01054-20UG / C01054-100UG
Package	5 µg / 20 µg / 100 µg
Description	<p>CD40, belonging to the TNF receptor family, is a cell surface protein generally expressed on B cells, dendritic cells, monocytes, thymic epithelial cells and on T cells at low levels. The membrane-anchored CD40 Ligand is expressed almost exclusively on activated CD4+ T lymphocytes. CD40 signaling plays a significant role immunoglobulin (Ig) class switching, and is critical for the proliferation and differentiation of B cells. A disease "immunodeficiency with hyper-IgM", characterized by failure to produce IgG, IgA and IgE due to failure expression of CD40L. The soluble form of CD40L is produced in vivo by an intracellular proteolytic processing of the entire TNF homologous region (18 kDa protein) of CD40L.</p>
Source	<i>Escherichia coli</i>
Sequence	<p>MQKGDQNPQIAAHVISEASSKTTSVLQWAEKGYTMSNNLVTLENGKQLTVKR QGLYYIYAQVTFCSNREASSQAPFIASLCLKSPGRFERILLRAANTHSSAKPCG QQSIHLGGVFELQPGASVFNVTDPQSQVSHGTGFTSFGLLKL with polyhistidine tag at the C-terminus</p>
Endotoxin level	<0.1 EU per 1 µg of the protein by the LAL method.
Activity	Measure by its ability to induce IL-8 secretion in human PBMCs. The ED ₅₀ for this effect is <5 ng/mL.
Purity	>95% as determined by SDS-PAGE.
Form	Lyophilized
Storage Buffer	Lyophilized from a 0.2 µm filtered solution of PBS, pH 8.0.
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile H ₂ O to a concentration not less than 200 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.
Stability & Storage	<p>This product is stable after storage at:</p> <ul style="list-style-type: none"> -20°C for 12 months in lyophilized state from date of receipt. -20°C or -80°C for 1 month under sterile conditions after reconstitution. <p>Avoid repeated freeze/thaw cycles.</p>



SDS-PAGE analysis of recombinant human CD40L

For research use only.