


**TRAIL (TNF-related apoptosis-inducing ligand), Human**

v. 231001

<b>Catalog number</b>	C01059-5UG / C01059-20UG / C01059-100UG
<b>Package</b>	5 µg / 20 µg / 100 µg
<b>Description</b>	TRAIL acts as a cytotoxic protein, through activating rapid apoptosis in tumor cells (not in normal cells). TRAIL conducts apoptosis through binding to DR4 and DR5, which are two death-signaling receptors belonging to the TNFR superfamily of transmembrane proteins. These receptor contain a cytoplasmic "death domain", which can initiate the cell's apoptotic process.
<b>Source</b>	<i>Escherichia coli</i>
<b>Sequence</b>	MRERGPQRVAAHITGTRGRSNTLSSPNSKNEKALGRKINSWESSRSGHSFLSN LHLRNGELVIHEKGFYYIYSQTYFRFQEEIKENTKNDKQMVQYIYKYTSYPDPILL MKSARNSCWSKDAEYGLYSIQGGIFELKENDRIFVSVTNEHLIDMDHEASFFG AFLVG with polyhistidine tag at the C-terminus
<b>Endotoxin level</b>	<0.1 EU per 1 µg of the protein by the LAL method.
<b>Activity</b>	Measure by its ability to induce cytotoxicity in L929 cells in the presence of actinomycin D. The ED <sub>50</sub> for this effect is 10.4-15.4 ng/mL.
<b>Purity</b>	>98% as determined by SDS-PAGE.
<b>Form</b>	Lyophilized
<b>Storage Buffer</b>	Lyophilized from a 0.2 µm filtered solution of PBS, pH 8.0.
<b>Reconstitution</b>	It is recommended to reconstitute the lyophilized protein in sterile H <sub>2</sub> O to a concentration not less than 200 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.
<b>Stability &amp; Storage</b>	This product is stable after storage at: <ul style="list-style-type: none"><li>• -20°C for 12 months in lyophilized state from date of receipt.</li><li>• -20°C or -80°C for 1 month under sterile conditions after reconstitution.</li></ul> Avoid repeated freeze/thaw cycles.



SDS-PAGE analysis of recombinant human TRAIL

*For research use only.*