


**FGF-23 (Fibroblast growth factor-23), Human**

v. 231001

<b>Catalog number</b>	C01113-5UG / C01113-20UG / C01113-100UG
<b>Package</b>	5 µg / 20 µg / 100 µg
<b>Description</b>	FGF-23 is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. FGF-23 is most highly expressed in bone, from which it circulates through the blood to regulate vitamin D and phosphate metabolism in kidney.
<b>Source</b>	<i>Escherichia coli</i>
<b>Sequence</b>	<p>           MYPNASPLLGSWSGGLIHLTYTATARNSTYHLQIHKNGHVDGAPHQTIYSALMIRS            EDAGFVVITGVMSRRYLCMDFRGNIFGSHYFDPENCRFQHQTLENGYDVYHS            PQYHFLVSLGRAKRAFLPGMNPYPYSQFLSRRNEIPLIHFNTPIPRRHTRSIED            DSERDPLNVLKPRARMTPAPASCSQELPSAEDNSPMASDPLGVVVRGGRVNTH            AGGTGPEGCRPFKFI with polyhistidine tag at the C-terminus         </p>
<b>Endotoxin level</b>	<0.1 EU per 1 µg of the protein by the LAL method.
<b>Activity</b>	Measure by its ability to induce proliferation in BaF3 cells transfected with human FGFR3c. The ED <sub>50</sub> for this effect is <0.3 µg/mL.
<b>Purity</b>	>98% as determined by SDS-PAGE.
<b>Form</b>	Lyophilized
<b>Storage Buffer</b>	Lyophilized from a 0.2 µm filtered solution of PBS, pH 8.0.
<b>Reconstitution</b>	It is recommended to reconstitute the lyophilized protein in sterile H <sub>2</sub> O to a concentration not less than 200 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.
<b>Stability &amp; Storage</b>	<p>This product is stable after storage at:</p> <ul style="list-style-type: none"> <li>-20°C for 12 months in lyophilized state from date of receipt.</li> <li>-20°C or -80°C for 1 month under sterile conditions after reconstitution.</li> </ul> <p>Avoid repeated freeze/thaw cycles.</p>



SDS-PAGE analysis of recombinant human FGF-23

---

*For research use only.*