

CXCL7 (C-X-C motif chemokine 7), Human

v. 231001

Catalog number	C01133-5UG / C01133-20UG / C01133-100UG
Package	5 µg / 20 µg / 100 µg
Description	The platelet-derived protein CXCL7 is a growth factor that belongs to the alpha-chemokine family. It is released in large amounts from platelets following their activation. It stimulates various processes including mitogenesis, synthesis of extracellular matrix, glucose metabolism and synthesis of plasminogen activator.
Source	<i>Escherichia coli</i>
Sequence	AELRCMCIKTTSGIHPKNIQSLEVIGKGTGHCNQVEVIATLKDGRKICLDPDAPRIK KIVQKKLAGDESAD with polyhistidine tag at the N-terminus
Endotoxin level	<0.1 EU per 1 µg of the protein by the LAL method.
Activity	Measure by its ability to chemoattract BaF3 cells transfected with human CXCR2. The ED ₅₀ for this effect is <0.5 ng/mL.
Purity	>98% as determined by SDS-PAGE.
Form	Lyophilized
Storage Buffer	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile H ₂ O to a concentration not less than 200 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.
Stability & Storage	This product is stable after storage at: <ul style="list-style-type: none">• -20°C for 12 months in lyophilized state from date of receipt.• -20°C or -80°C for 1 month under sterile conditions after reconstitution. Avoid repeated freeze/thaw cycles.



SDS-PAGE analysis of recombinant human CXCL7

For research use only.