


IL-24 (Interleukin-24), Mouse

v. 231101

Catalog number	C02028-5UG / C02028-20UG / C02028-100UG
Package	5 µg / 20 µg / 100 µg
Description	<p>Interleukin-24 (IL-24) is a cytokine belonging to the IL-10 family of cytokines that signals through two heterodimeric receptors: IL-20R1/IL-20R2 and IL-22R1/IL-20R2. This interleukin is also known as melanoma differentiation-associated 7 (mda-7) due to its discovery as a tumour suppressing protein. IL-24 appears to control in cell survival and proliferation by inducing rapid activation of particular transcription factors called STAT1 and STAT3. This cytokine is predominantly released by activated monocytes, macrophages and T helper 2 (Th2) cells and acts on non-haematopoietic tissues such as skin, lung and reproductive tissues. IL-24 performs important roles in wound healing, arthritis, psoriasis and cancer. Several studies have shown that cell death occurs in cancer cells/cell lines following exposure to IL-24. The gene for IL-24 is located on chromosome 1 in humans.</p>
Source	<i>Escherichia coli</i>
Sequence	<p>MQEFRFGSCQVTGVVLPPELWEAFWTVKNTVQQTQDDITSIRLLKPQVLRNVSGA ESCYLAHSLKLFYLN TVFKNYHSKI AKFKVLR SFSTLANNFIVIMSQLQPSKDNS MLPIS ESAHQRFLLFRRAFKQLDTEVALVKAFGEVDILLTWMQKFYHL with polyhistidine tag at the C-terminus</p>
Endotoxin level	<0.1 EU per 1 µg of the protein by the LAL method.
Activity	Measure by its ability to induce proliferation in BaF3 cells transfected with human IL-20 R alpha and human IL-20 R beta. The ED ₅₀ for this effect is <0.3 ng/mL.
Purity	>95% as determined by SDS-PAGE.
Form	Lyophilized
Storage Buffer	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile H ₂ O to a concentration not less than 200 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.
Stability & Storage	<p>This product is stable after storage at:</p> <ul style="list-style-type: none"> -20°C for 12 months in lyophilized state from date of receipt. -20°C or -80°C for 1 month under sterile conditions after reconstitution. <p>Avoid repeated freeze/thaw cycles.</p>



SDS-PAGE analysis of recombinant mouse IL-24

For Research Use Only.